

Contact Infection Control prior to discontinuing Precautions

Washington State Hospital Association



Salmon HEX F9D7D4 Last revised 3/03/20

Special Droplet/Contact Precautions

If patient has diarrhea and/or C. difficle add Contact Enteric Precautions

LIMIT VISITORS AND DOCUMENT ANYONE THAT HAS CONTACT WITH THE PATIENT AND NOTIFY EMPLOYEE HEALTH

Display sign outside the door. At patient discharge, remove sign <u>after</u> room is terminally cleaned.

For use with:

Respiratory viruses including COVID-19

Dishes/Utensils:

No special precautions. Kitchenware sanitized in dishwasher.

Equipment and Supplies:

- Only essential supplies in room.
- Use dedicated or disposable equipment when available.
- Minimize use of cellphones/pagers.
- Clean and disinfect reusable equipment including intravenous pumps, cell phone or pagers (if used in room), and other electronics, supplies, and other equipment prior to removing from patient's room.
- Ensure blood pressure cuff and stethoscope are cleaned and disinfected between patients.

Waste and Linen Management:

For COVID-19, follow local and state public health guidelines Category B for medical waste handling. Bag linen in the patient's room.

Private Room:

If not available, room with patients that have the same organism but no other infection.

Room Cleaning:

Routine cleaning procedures with addition of cubical curtain changes per hospital procedure.

Transport:

Essential transport only. Patient should remain in room except for medical necessity. Patient should wash their hands. Place patient in clean gown. Place surgical mask on patient. Clean and disinfect transport vehicle. Alert receiving department regarding patient's isolation precaution status.

Personal Protective Equipment:

